

TWO NEW SPECIES OF BRAZILIAN PHILODENDRON (ARACEAE)

George S. Bunting

Philodendron kautskyi Bunt., sp. nov.

Herba scandens. Caulis 1.8 cm diam. internodiis ad 2 cm longis. Petiolus teres 19-23.5 cm longus 0.8 cm crassus (in medio), parte vaginata 5-6 cm longa ala altera 0.8 cm lata. Foliorum lamina subcoriacea ambitu late ovata vel cordiformis 25.5 cm longa 22.5 cm lata (in loco 1 cm supra basim costae), ad apicem obtusa abrupte et breviter acuminata, ad basim profunde cordata lobis posticis rotundatis sinu parabolico 9 cm longo sejunctis, adaxiale nitida intense viridis costa convexa, abaxiale vix nitida, nervis lateribus 1. utrinque 4 sub angulo ca. 55° abeuntibus, uterque costula in sinu per 1.5-2 cm nuda nervis principalibus latere exteriore 2. Inflorescentia solitaria [specimine ante anthesin]. Pedunculus 4 cm longus. Spatha 12.5 cm longa (quam spadix 1.2 cm longiora), extra ubique viridis, intra viridula, tubo plus minusve fusiformi. Flores pistillati 3.5 cm longi, stylo gracili ovarium globoideum paullum superanti, stigmate globoso, ovario 8-10-loculari, ovulis in quoque loculo 5 (4-6) basim versus centraliter affixis.

TYPUS: George S. Bunting B82-29. Domingos Martins, Espírito Santo, BRAZIL, ca. 800 m. February, 1982 [Holotypus: NY]. The specimen was collected by Mr. Roberto Kautsky and brought to me in Rio de Janeiro by Mr. Luiz Knud Correia de Araujo; both gentlemen are dedicated plantsmen.

Although *P. kautskyi* clearly pertains to the subgenus *Philodendron*, the ovary characters do not correspond well to any of the sections of the genus as circumscribed by Krause (Pflanzenreich [Engler]. Heft 60. 1913).

Philodendron spiritus-sancti Bunt., sp. nov.

Herba scandens. Caulis 60 cm vel ultra longus 4.5 cm diam. internodiis brevissimis (0.5 cm) attamen cicatricibus foliorum delapsorum ca. 1.8 cm diam. Petiolus abaxiale rotundatus, adaxiale late canaliculatus, apicem versus semiteres adaxiale plus minusve conexus, 53 cm longus 0.9 cm crassus (in medio), sordide luteo-viridis multis maculis ellipticis vinosis et abaxiale vinaceo-suffusus, per 10 cm superiores atrovinosus, parte vaginata 9-11 cm longa alis ochraceis. Foliorum lamina subcoriacea, plana attamen margine revoluto et vino (1 mm), ambitu longe triangularis basim versus lobii antici aliquantum repanda, ad apicem acuminata (1-1.5 cm) ad basim sagittata, usque 80 cm longa trans apicem loborum posticorum 16 cm lata, trans insertionem petioli 15 cm lata, in medio lobi antici 12 cm lata; lobi postici oblongo-longe triangulares usque 15.5 cm longi 4.7 cm lati et 18 cm longi 4.3 cm lati (in folio eodem) sinu longe trianguli 15 cm longo sejuncti; abaxiale ubique verna ner-




Fig. A-D, *Philodendron kautskyi* Bunt. A, leaf and inflorescence, much reduced; B, cross-section of petiole at midpoint, ca. 1/2; C, pistil, enlarged; D, ovules from a single locule, much enlarged. Fig. E-H, *P. spiritus-sancti* Bunt. E, leaf and inflorescence, much reduced; F, cross-section of petiole at midpoint, ca. 1/2; G, pistil, enlarged; H, ovules from a single locule, much enlarged.

vatione vinosa obscuriore, costa valde convexa; adaxiale impolita intense viridis nervatione flavo-virenti, costa plana vel vix convexa 7 mm lata, nervis lateralibus 1. sulcatis utrinque 5-6 sub angulo 45°-65° abeuntibus, uterque costula in sinu per 5-6 cm nuda et in parte distali ex margine tantum 8-11 mm remota nervis principali bus latere exteriore 3. Inflorescentiae solitariae vel binae. Pedunculus 6 cm longus viridis ad apicem vinosus saturatus. Spatha 16.5 cm longa, non constricta et ubique plus minusve eodem diametro (1.9 cm), extra viridis maculis ellipticis vinosis intra viridi-cremea sursum eburnea, late aperiens. Spadix spatham aequans insertione obliquissima, 14.5 cm longus parte pistillata 6 (2.4) cm longa 1.3 cm crassa, parte staminata sterili 1.9(1.7) cm longa 1.4 cm crassa (ad apicem 1.1 cm crassa), fertili 1.2-1.3 cm crassa. Flores pistillati cylindrici stigmate coronato, 3.8 cm longi 1.8-2 mm crassi, ovario 10-11(-13)-loculari, ovulis in quoque loculo 3 (-4) basim versus centraliter affixis.

TYPUS: George S. Bunting B82-28. In rainforest, "Morro do Suido", Domingos Martins, Espírito Santo, BRAZIL, ca. 800 m. January 20, 1982 [Holotypus: NY]. This specimen was also collected by Mr. Roberto Kautsky and brought to me in Rio de Janeiro by Mr. Luiz Knud Correia de Araujo.

Philodendron spiritus-sancti pertains to sect. Oligospermium Engl. § Macrolbelium Schott.

ERRATA

re BUNTING, G. S., New taxa of Venezuelan Araceae. Phytologia 60: 293-344. 1986.

page	line	reads:	should read:
311	27	en km 25 del trayecto al este de San Cristóbal	entre km 11 y 25 al este de San Cristóbal, o sea 0-14 km al este del Chorro del Indio
311	28	ca. 1100 m, 7 de junio, 1973	1100-1150 m, 24 de abril, 1976
332	4	en km 25 al este de San Cristóbal	entre km 11 y 25 al este de San Cristóbal, o sea 0-14 km al este del Chorro del Indio
332	5	ca. 1100 m, 7 de junio, 1973	1100-1150 m, 24 de abril, 1976
340	scale	10 mm	10 cm